

TABLE OF CONTENTS

CHAPTER I - KNOWN ALGORITHM FOR REAL NUMBERS	11
Section 1.0. Introduction	11
Section 1.1. The Euclidean Algorithm (EA), Continued Fraction Algorithm and Euler-Lagrange Theorem (2-ELT) for Quadratics	12
Section 1.2. The Euclidean Algorithm is the Euler System for Quadratics	17
Section 1.3. Jacobi Algorithm (JA)	18
Section 1.4. Perron Algorithm (PA), and the General Continued Fraction Algorithm	21
Section 1.5. Hasse and Bernstein Algorithm (HBA)	22
 CHAPTER II – BAICA’S GEBNERALIZED EUCLIDEAN ALGORITHM (BGEA)	 24
Section 2.0. Introduction	24
Section 2.1. Baica’s General Euclidean Algorithm (BGEA) a different proof from the initial one for the necessary condition of its restricted periodicity	24
2.1* A periodic (BGEA); $n = 4$	28
2.2* A periodic (BGEA); $n = 5$	35
2.3* The (BGEA) for the general case $n > 2$	39
Section 2.2. The proof of the sufficient condition for the restricted periodicity of (BGEA).	49
Section 2.3. (BGEA) an n -dimensional equivalent of Euler – Lagrange theorem (n -ELT).	50
Section 2.4. (BGEA) is the Euler System for the Algebraic Number Theory in n -dimensions.	51
 CHAPTER III - THE ORIGINAL FERMAT’S LAST THEOREM STATED IN EUCLIDEAN TERMS (EFLT) AND THE ELLIPTICAL FERMAT’S LAST THEOREM (ELFLT).	 53
Section 3.0. Introduction.	53
Section 3.1. The Euclidean character of the original Fermat’s Last Theorem (EFLT).	53
Section 3.2. Explicit solutions for Hasse’s diophantine equations $a^2 \pm a b + b^2 = c^2$	54
3.1* Definitions and previous results.	55
3.2* Euler-Pell’s equation: $u^2 - 3v^2 = 1$, $n = 0, 1$,	57
3.3* A solution of $a^2 + a b + b^2 = c^2$	58
3.4* Solution of $a^2 - a b + b^2 = c^2$	60
3.5* Second solution of $a^2 - a b + b^2 = c^2$	63
3.6* Perimeters and areas.	65
3.7* Entirely new RPT-S.	65
3.8* The main result.	68

TABLE OF CONTENTS

Section 3.3. Units from (BGEA) play an important role in solving (EFLT). 70

Section 3.4. (BGEA) restricted periodicity and the Euclidean Solution of the original Fermat's Last Theorem stated in Euclidean terms (EFLT). 71

Section 3.5. More explanations about Baica's proof of the Euclidean Fermat's Last Theorem (EFLT). 75

Section 3.6. A Wiles's attempt to prove the Elliptical Fermat's Last Theorem (ELFLT). 81

Section 3.7. G. Faltings finishes the proof of the Elliptical Fermat's Last Theorem (ELFLT). 82

Section 3.8. Elliptical Fermat's Last Theorem (ELFLT) proved by Faltings is equivalent to (not the same as) the original Fermat's Last Theorem (EFLT) stated by Fermat in Euclidean terms and proved by Baica. 83

Section 3.9. Conclusions. 84

REFERENCES 87

AFTERWORD 91

CHAPTER III - THE ORIGINAL FERMAT'S LAST THEOREM
STATED IN EUCLIDEAN TERMS (EFLT) AND THE
ELLIPTICAL FERMAT'S LAST THEOREM

Section 3.1. The Euclidean character of the original Fermat's Last Theorem (EFLT). 70

Section 3.2. Elliptic solutions for Fermat's diophantine equations $x^n + y^n = z^n$ 71

3.2* Definitions and previous results. 72

3.2* Euler-Pell's equation $x^2 - Dy^2 = 1$, $x = 2, y = 1$ 73

3.2* A solution of $x^2 + 5y^2 = z^2$ 74

3.2* Solution of $x^2 - 5y^2 = z^2$ 75

3.2* Special solution of $x^2 - 5y^2 = z^2$ 76

3.2* Fermat's last theorem. 77

3.2* Fermat's last theorem. 78

**THE ALGORITHMIC SOLUTION
OF THE ORIGINAL
EUCLIDEAN FERMAT'S LAST THEOREM**

Author

Prof.Dr. Malvina BAICA
UNIVERSITY OF WISCONSIN – WHITEWATER
U. S. A.