

Italian Journal of Pure and Applied Mathematics

estratto

EDITOR-IN-CHIEF


Piergiulio Corsini

EDITORIAL BOARD

Luisa Arlotti
Malvina Baica
Adriano Barlotti
Federico Bartolozzi
Carlo Cecchini
Stephen Comer
Gianni Dal Maso
Bal Kishan Dass
Giacomo Della Riccia
Franco Eugeni
Giovanni Falcone
Antonino Giambruno
Furio Honsell

James Jantosciak
Jaroslav Ježek
Tomas Kepka
David Kinderlehrer
Andrzej Lasota
Violeta Leoreanu
Mario Marchi
Angelo Marzollo
Jean Mittas
Chin-Hong Park
Rodolfo Permutti
Livio C. Piccinini
Goffredo Pieroni

Ivo Rosenberg
Maria Scafati Tallini
Alessandro Silva
Sergio Spagnolo
Yves Sureau
Jeno Szepe
Carlo Tasso
Ioan Tofan
Hans Weber
Fabio Zanolin
Paolo Zellini



FORUM

SOLUTION OF GOLDBACH'S CONJECTURE

Malvina Baica

*University of Wisconsin
Whitewater, Wisconsin 53190
U.S.A.*

Abstract. An exact expression is derived from the Hardy–Littlewood function $\nu(t)$ by means of the Laplace transform. The application of the generalized final value theorem allows us to prove a result that essentially is the hypothetical asymptotic formula for the Hardy–Littlewood function $\nu(t)$, with corresponding remainder term. The final step of this paper is to evaluate the difference between the exact and the asymptotic value, in order to prove the validity of the hypothesis.

2000 Mathematics Subject Classification: 11A, 11Z05, 11R04

Keywords: Hardy–Littlewood function, Laplace transform.

0. Introduction

Goldbach's conjecture is an open problem from 1742, when Christian Goldbach, in a letter to Euler enunciated it as a conjecture that "every even number greater than two can be expressed as the sum of two primes".

Aldo Peretti wrote several papers about this subject published in the *Bulletin of Number Theory* (B.N.T.) [5] and in *Notes in Number Theory and Discrete Mathematics* (N.N.T.D.M.) [6], [7], [8].

In order to prove the Goldbach hypothesis, it is necessary to proceed in two steps.

First, to prove the Hardy–Littlewood hypothetical asymptotic formula, which was proved by Peretti in his 1997 paper [8].

Peretti [8] used an incorrect formula

$$(0.1) \quad \nu(t) = L^{-1}\{e^s(1 - e^{-s})^2 L^2(\vartheta(u))\}$$

for $\nu(t)$, instead of

$$(0.2) \quad \nu(t) = L^{-1}\{(1 - e^{-s})^2 L^2(\vartheta(u))\}$$

or

$$\nu(t) = L^{-1}\{L^2(\Delta\vartheta(u))\}$$

which we found to be correct.

However, Peretti's proof for (0.1) is valid for (0.2).

Also, we found an incorrect statement in theorem (B) of [8], which we should state as follows:

$$\begin{aligned} \text{If } \quad & F(t) \sim G(t) \quad \text{when } t \rightarrow \infty, \text{ then} \\ & f(s) \sim g(s) \quad \text{when } s \rightarrow 0, \text{ where} \\ & f(s) = L\{F(t)\} \quad \text{and } g(s) = L\{G(t)\} \end{aligned}$$

and, conversely, if some Tauberian condition is fulfilled.

In **step 2** we determine the error of the Hardy–Littlewood formula. This is the most difficult part and cannot be done by the circle–method.

Thus, we use the Laplace transform as a principal and crucial tool, which makes it possible to solve this problem.

As we can see, Peretti’s long and hard work on this problem with some corrections and some final modifications brought us to the final proof of this very challenging problem.

1. Derivation of a basic formula

Using the Chebishev function $\vartheta([u])$ defined by

$$(1.1) \quad \vartheta([u]) = \sum_{p \leq [u]} \log p$$

where p denotes the prime numbers, we have:

$$(1.2) \quad \Delta\vartheta([u]) = \vartheta([u]) - \vartheta([u-1]) = \begin{cases} \log p & \text{if } [u] = p \\ 0 & \text{in any other case} \end{cases}$$

Hence, if we form

$$(1.3) \quad \nu(t) = \sum \Delta\vartheta([u_1]) \sum \Delta\vartheta([u_2])$$

where the sums are extended to those values of u_1 and u_2 such that

$$t = u_1 + u_2$$

evidently, $\nu(t)$ is the Hardy–Littlewood function

$$(1.4) \quad \nu(t) = \sum_{p_1 + p_2 = t} \log p_1 \log p_2.$$

Immediately, from (1.3) we can see that we also have:

$$(1.5) \quad \nu(t) = \sum_{u_1 < t} \Delta\vartheta([u_1]) \sum_{u_2 < t - u_1} \Delta\vartheta([u_2])$$

or

$$(1.6) \quad \nu(t) = \sum_{u_1 < t} \Delta\vartheta([u_1]) \sum_{u_1 < t} \Delta\vartheta([t - u_1]).$$

In fact, in the first sum of (1.5) we choose arbitrarily any integer $u_1 < t$; while in the second we are forced to restrict the sum to the $u_2 < t - u_1$.

As $\vartheta(u)$ is a step function the sums on the right-hand side of (1.6) are also step functions, that can be replaced by the corresponding integrals, so that we can write:

$$(1.7) \quad \nu(t) = \int_0^x \Delta\vartheta([u])du \int_0^x \Delta\vartheta([t-u])du.$$

Here, the question of what value x must assume, has to be examined.

We have that

$$(1.8) \quad \int_a^{a+1} \Delta\vartheta([u])du = \begin{cases} \log a & \text{if } a = p \\ 0 & \text{if } a \neq p \end{cases}$$

Hence it is useless, to put $x = t + 1$ in (1.7) because t cannot be an odd prime, as it is assumed to be an even number. In conclusion, in (1.7) we can choose $x = t$ so that

$$(1.9) \quad \nu(t) = \int_0^t \Delta\vartheta([u])du \int_0^t \Delta\vartheta([t-u])du = \int_0^t \Delta\vartheta([u_1])du_1 \int_0^{t-u_1} \Delta\vartheta([u_2])du_2.$$

(This is the integral version of (1.5).)

Evidently, the right hand member is, by definition, the convolution of the two functions

$$(1.10) \quad \nu(t) = \Delta\vartheta([u]) * \Delta\vartheta([u])$$

with current notation.

Consequently, if we take the Laplace transform L of both sides, by usual rules, we obtain

$$(11) \quad \begin{aligned} L\{\nu(t)\} &= L\{\Delta\vartheta([u])\}L\{\Delta\vartheta([u])\} = L^2\{\Delta\vartheta([u])\} \\ L\{\nu(t)\} &= L^2\{\Delta\vartheta([u])\} \end{aligned}$$

and this is our basic formula needed.

2. The evaluation of $L\{\Delta\vartheta([u])\}$

According to [9], vol.4, p.2, formula 8, we have:

$$(2.1) \quad L\{\vartheta([u])\} = \frac{1 - e^{-s}}{s} \sum_{k=0}^{\infty} \vartheta(k)e^{-ks}.$$

From this it follows that

$$(2.2) \quad L\{\Delta\vartheta([u])\} = \frac{1 - e^{-s}}{s} \sum_p \log p e^{-ps}$$

when we take into account (1.2).

3. The poles of $f(s) = \sum_{\mathbf{p}} \log \mathbf{p} e^{-\mathbf{p}s}$

Setting $e^{-s} = x$ we have:

$$(3.1) \quad \sum_{\mathbf{p}} \log \mathbf{p} e^{-\mathbf{p}s} = \sum_{\mathbf{p}} \log \mathbf{p} x^{\mathbf{p}} = F(x).$$

It is known that $F(x)$ has infinitely many simple poles on the circle $|x| = 1$. Hence, $f(s)$ has infinitely many poles on the line $\sigma = \operatorname{Re}(s) = 0$.

For further calculation using with the Laplace transforms, we are interested in the residues of these poles at $x = e^{2\pi i h/q}$ as a primitive root of unity of order q . Consequently, we must assume that $(h, q) = 1$.

The answer is given by Theorem 248, Ch.5, p. 212 of [4]

$$(3.2) \quad F(x) \sim \frac{\mu(q)}{\varphi(q)(1 - x e^{-2\pi i h/q})} = -\frac{\mu(q)e^{2\pi i h/q}}{\varphi(q)(x - e^{2\pi i h/q})}$$

where $\mu(q)$ (is Möbius function) $= (-1)^r$ if q has r factors and $\varphi(q)$ (is Euler's function) is the number of integers $\leq q$ and coprime with q .

More exactly, as h can take the values $h = 0, 1, 2, \dots, q-1$ with the restriction $(h, q) = 1$, we can write that $F(x)$, in the surroundings of the $\varphi(q)$ primitive roots of unity of order q , is as

$$(3.3) \quad F(x) \sim -\sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)e^{2\pi i h/q}}{\varphi(q)(x - e^{2\pi i h/q})}.$$

When we extend this reasoning to all the roots of order $q \leq N$, we get the approximation

$$(3.4) \quad F(x) \sim -\sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)e^{2\pi i h/q}}{\varphi(q)(x - e^{2\pi i h/q})}.$$

If we desire an error term for this approximation, we can use the remainder term given by Theorem 248 of [4]. However, for our purposes it is sufficient to put in evidence the number and the kind of the poles of $F(x)$ only.

Next if we put $x = e^{-s}$ in (3.4), we obtain:

$$(3.5) \quad F(e^{-s}) = f(s) \sim -\sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)e^{2\pi i h/q}}{\varphi(q)(e^{-s} - e^{2\pi i h/q})}.$$

Again here we are interested in the calculation of the poles of $f(s)$, and in the value of their residues.

We have that the residue of

$$g(s) = \frac{1}{e^{-s} - e^{2\pi i h/q}} \quad .$$

at the simple pole $s = -2\pi ih/q$ is

$$-e^{-2\pi ih/q}.$$

Thus we can write (3.5) alternatively as

$$(3.6) \quad f(s) \sim \sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)}{\varphi(q)(s + 2\pi ih/q)}.$$

The right hand side of the formula (3.6) is known as the Farey dissection of order N of $f(s)$.

Replacing (3.6) in (2.2) we get:

$$(3.7) \quad L\{\Delta\vartheta([u])\} \sim \frac{1 - e^{-s}}{s} \sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)}{\varphi(q)(s + 2\pi ih/q)}$$

and taking into account formula (1.11) we have:

$$(3.8) \quad L\{\nu(t)\} \sim \left(\frac{1 - e^{-s}}{s}\right)^2 \left\{ \sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu(q)}{\varphi(q)(s + 2\pi ih/q)} \right\}^2.$$

If we wish to pass from the asymptotic sign to the equality sign, we must write:

$$(3.9) \quad L\{\nu(t)\} = \left(\frac{1 - e^{-s}}{s}\right)^2 \left\{ \lim_{N \rightarrow \infty} B_N^2(s) \right\}$$

where $B_N(s)$ is given in the second bracket on the right hand side of (3.8). The importance of formula (3.8) lies in the fact that it exhibits the poles of $L\{\nu(t)\}$, which is the basis for the evaluation of $\nu(t)$ by the residues method.

4. Abelian and Tauberian theorems concerning the Laplace transform

Suppose that

$$f(s) = L\{F(u)\} \quad \text{and} \quad g(s) = L\{G(u)\}$$

and

$$(4.1) \quad f(s) \sim g(s) \quad \text{when} \quad s \rightarrow 0^+ \quad \text{means that} \quad \lim_{s \rightarrow 0} f(s)/g(s) = 1$$

and similarly

$$(4.2) \quad F(t) \sim G(t) \quad \text{when} \quad t \rightarrow \infty \quad \text{means that} \quad \lim_{t \rightarrow \infty} F(t)/G(t) = 1.$$

An implication from (4.2) to (4.1) is valid under rather general assumptions (that are fulfilled in our case), and is called an "Abelian theorem".

In the theory of the Laplace transform it is called the generalized theorem of the final value, which also is valid even for discontinuous functions.

The converse implication from (4.1) to (4.2) is true under additional assumptions only, and it is named a "Tauberian Theorem", which is available a wide variety of cases.

However, we shall not appeal to such kind of theorems.

5. Application of the Abelian theorem to (3.8)

Let

$$\nu^*(t) = \lim_{t \rightarrow \infty} \nu(t)$$

be such that

$$\nu^*(t) \sim \nu(t).$$

Then, by the Abelian theorem, we get

$$L\{\nu^*(t)\} = \lim_{t \rightarrow \infty} L\{\nu(t)\} = \lim_{s \rightarrow 0} \left(\frac{1 - e^{-s}}{s} \right)^2 B_N^2(s)$$

where $B_N(s)$ is as defined before in (3.8).

This reduces to:

$$(5.1) \quad L\{\nu^*(t)\} = B_N^2(s)$$

from which it follows

$$(5.2) \quad \nu^*(t) = L^{-1}\{B_N^2(s)\}.$$

More exactly, the passage from (5.1) to (5.2) should be written as

$$(5.3) \quad \nu_m^*(t) = \frac{\nu^*(t+0) + \nu^*(t-0)}{2} = L^{-1}\{B_N^2(s)\}$$

because $\nu^*(t)$ is a discontinuous function.

6. Evaluation of $\nu_m^*(t)$ according to (5.3)

Due to a well-known theorem [12], we have:

$$(6.1) \quad \nu_m^*(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} B_N^2(s) e^{ts} ds$$

where the contour between $c \pm i\infty$ is a Bromwich contour, that contains all the poles of $B_N^2(s)$.

In the usual way, it is evaluated as the sum of the residues of the integrand at their poles, within the Bromwich contour.

Now, we have that:

$$\begin{aligned}
 B_N^2(s) &= \sum_{q=1} \sum_{h=0, (h,q)=1} \frac{\mu^2(q)}{\varphi^2(q)(s + 2\pi i h/q)^2} \\
 (6.2) \quad &+ \sum_{q_1=1}^N \sum_{h_1=0, (h_1,q_1)=1}^{q_1-1} \sum_{q_2=1}^N \sum_{h_2=0, (h_2,q_2)=1}^{q_2-1} \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)\varphi(q_2)(s+2\pi i h_1/q_1)(s+2\pi i h_2/q_2)} = \\
 &= \sum_{q=1}^N \sum_{h=0, (h,q)=1}^{q-1} \frac{\mu^2(q)}{\varphi^2(q)(s + A)^2} + \\
 &+ \sum \sum \sum \sum \sum \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)\varphi(q_2)} \frac{1}{(A_2 - A_1)} \left\{ \frac{1}{s + A_1} - \frac{1}{s + A_2} \right\}.
 \end{aligned}$$

with the obvious notation for A, A_1, A_2 and the sums.

The residues of the integrand at the double poles $1/(s + A)^2$ are easily found to be

$$\frac{\mu^2(q)}{\varphi^2(q)} t e^{-At}.$$

The residue of the integrand at the simple pole $1/(s + A_1)$ is found to be

$$\frac{\mu(q_1)}{\varphi(q_1)} e^{-A_1 t}.$$

Hence, collecting partial results, we have:

$$\begin{aligned}
 (6.3) \quad \nu_m^*(t) &= \sum \sum \frac{\mu^2(q)}{\varphi^2(q)} t \cdot e^{-At} + \\
 &+ \sum \sum \sum \sum \sum \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)\varphi(q_2)} \frac{e^{-A_1 t} - e^{-A_2 t}}{A_2 - A_1}.
 \end{aligned}$$

In a more simplified way, we can argue alternatively that the inverse Laplace transform of the finite sums in (6.2) is the sum of the inverse Laplace transform of its terms, and after that we appeal to the tables.

The quadruple sum in (6.3) can be maximized as follows. The minimal value of $A_2 - A_1$ is $2\pi i(h_1/q_1 - h_2/q_2)$ where both fractions are contiguous Farey fractions.

It is known [10] that

$$\min \left(\frac{h_1}{q_1} - \frac{h_2}{q_2} \right) = \frac{1}{N(N-1)}$$

so that

$$\left| \frac{1}{A_2 - A_1} \right| \leq \frac{N(N-1)}{2\pi}.$$

Furthermore

$$|e^{-A_1 t} - e^{-A_2 t}| \leq 2.$$

Hence:

$$\left| \sum_{h_1=0, (h_1, q_1)=1}^{q_1-1} \frac{e^{-A_1 t} - e^{-A_2 t}}{A_2 - A_1} \right| < \sum_{\substack{h_1=0, (h, q)=1 \\ (h_1, q_1)=1}}^{q_1-1} \left| \frac{e^{-A_1 t} - e^{-A_2 t}}{A_2 - A_1} \right| \leq \frac{\varphi(q_1)N(N-1)}{\pi}$$

with a similar result for the sum along h_2 .

It follows from the preceding inequalities:

$$\begin{aligned} \left| \sum_{q_1} \sum_{h_1} \sum_{q_2} \sum_{h_2} \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)\varphi(q_2)} \frac{e^{-A_1 t} - e^{-A_2 t}}{A_2 - A_1} \right| &< \sum_{q_1=1}^N \sum_{q_2=1}^N \frac{1}{\pi^2} N^2 (N-1)^2 = \\ &= \frac{1}{\pi^2} N^4 (N-1)^2 < \frac{N^6}{\pi^2} \end{aligned}$$

so that (6.3) can be written as:

$$(6.4) \quad \nu_m^*(t) = \sum_{q=1}^N \sum_{h=0, (h, q)=1}^{q-1} \frac{\mu^2(q)}{\varphi^2(q)} e^{-At} + \delta \frac{N^6}{\pi^2}$$

where $-1 < \delta < 1$. But

$$\sum_{h=0, (h, q)=1}^{q-1} e^{-At}$$

is Ramanujan's sum $Cq(t)$. Hence (6.4) can be written as:

$$(6.5) \quad \nu_m^*(t) = \sum_{q=1}^N \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) \cdot t + \delta_1 \frac{N^6}{\pi^2}$$

where $-1 < \delta_1 < 1$.

7. The evaluation of $\nu_m^*(t)$ and $f_N(t)$

Once we have evaluated

$$\nu_m^*(t) = L^2\{B_N(s)\}$$

it will be easy to evaluate

$$(7.1) \quad f_N(t) = L^{-1} \left\{ \left(\frac{1 - e^{-s}}{s} \right) B_N(s) \right\}^2$$

whose connection with (3.9) is evident.

Let

$$L^{-1}(f_1(s)) = F_1(t) \text{ and } L^{-1}(f_2(s)) = F_2(t)$$

where

$$(7.2) \quad f_1(s) = \frac{(1 - e^{-s})^2}{s^2} \text{ and } f_2(s) = B_N^2(s).$$

Then $F_2(t) = \nu_m^*(t)$, as was seen before.

In regard to

$$f_1(s) = \left(\frac{1 - e^{-s}}{s} \right)^2$$

and from the tables of [9], Vol.5, p.59, formula (15), we obtain:

$$(7.3) \quad F_1(t) = L^{-1} \left\{ \frac{(1 - e^{-s})}{s^2} \right\} = \begin{cases} t & \text{if } 0 < t < 1 \\ 2 - t & \text{if } 1 < t < 2 \\ 0 & \text{otherwise} \end{cases}$$

By the convolution theorem for the inverse Laplace transform we have:

$$f_N(t) = L^{-1} \{ f_1(s) \cdot f_2(s) \} = \int_0^t F_1(u) F_2(t - u) du.$$

Replacing the values given by (7.2) and (7.3) in the above transform we obtain:

$$(7.4) \quad f_N(t) = \int_0^t F_1(u) \nu_m^*(t - u) du = \int_0^1 u \nu_m^*(t - u) du + \int_1^2 (2 - u) \nu_m^*(t - u) du.$$

8. Finding the limit of $\nu_m^*(t)$ when $N \rightarrow \infty$

Now we find:

$$(8.1) \quad \eta(t) = \lim_{N \rightarrow \infty} \nu_m^*(t).$$

For this we need:

Lemma 1. This lemma proves:

$$(L.1) \quad \left| \sum_{q=N+1}^{\infty} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) \right| < 10240 \sqrt[3]{t} \frac{\log^3 N}{N}.$$

Proof of Lemma 1. According to [13], p.35, formula (3.23), we have:

$$(L.2) \quad \sum_{q>N} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) = \sum_{d|t} \frac{\mu^2(d)}{\varphi(d)} \sum_{N/d < q, (q,t)=1} \frac{\mu(q)}{\varphi^2(q)}.$$

Now

$$(L.3) \quad \sum_{N/d < q, (q,t)=1} \left| \frac{\mu(q)}{\varphi^2(q)} \right| < \sum_{N/d < q, (q,t)=1} \frac{1}{\varphi^2(q)} < \sum_{N/d < q} \frac{1}{\varphi^2(q)}.$$

But

$$(L.4) \quad \varphi(n) > \frac{n}{2 + \log n} > \frac{n}{8 \log n}$$

for every $n > 1$ [2], so that

$$(L.5) \quad \sum_{N/d < q} \frac{1}{\varphi^2(q)} < 64 \sum_{N/d < q} \frac{\log^2 q}{q^2} < 64 \int_{N/d-1}^{\infty} \frac{\log^2 u}{u^2} du$$

and

$$\int_Q^{\infty} \frac{\log^2 u}{u^2} du = \frac{\log^2 Q}{Q} + \frac{2 \log Q}{Q} < 3 \frac{\log^2 Q}{Q} \quad \text{if } Q > e^2.$$

Hence

$$(L.6) \quad \begin{aligned} \int_{N/d-1}^{\infty} \frac{\log^2 u}{u^2} du &= \int_{N/d-1}^{N/d} \frac{\log^2 u}{N/d} du + \int_{N/d}^{\infty} \frac{\log^2 u}{u^2} du < \\ &< \frac{\log^2(N/d - \vartheta)}{(N/d - \vartheta)^2} + 3 \frac{\log^2(N/d)}{N/d} < 5 \frac{\log^2(N/d)}{N/d} \end{aligned}$$

where $0 < \vartheta < 1$ and the last inequality holds for every N/d .

Replacing it in (L.5) yields:

$$(L.7) \quad \sum_{N/d < q} \frac{1}{\varphi^2(q)} < 320 \frac{\log^2(N/d)}{N/d}.$$

From (L.2), (L.3), and (L.7) follows:

$$(L.8) \quad \begin{aligned} \left| \sum_{N < q} \frac{\mu^2(d)}{\varphi^2(q)} Cq(t) \right| &< \sum_{d|t} \frac{\mu^2(d)}{\varphi(d)} 320 \frac{\log^2(N/d)}{N/d} < \\ &< 320 \sum_{d|t} \frac{8 \log d}{d} \frac{\log^2(N/d)}{N/d} = 2560 \sum_{d|t} \frac{\log d \log^2(N/d)}{N} < \\ &< 2560 \sum_{d|t} \frac{\log^3(N/d)}{N} < 2560 d(t) \frac{\log^3 N}{N} \end{aligned}$$

where $d(t)$ denotes the quantity of divisors of t .

According to [3], holds that

$$(L.9) \quad d(t) < 4 \sqrt[3]{t}.$$

Replacing this value in (L.8), we obtain (L.1).

Combining (6.5) and Lemma 1, we get

$$(8.2) \quad \nu_m^*(t) = \sum_{q=1}^{\infty} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) \cdot t + \delta_1 \frac{N^6}{\pi^2} + \delta_2 10240 \sqrt[3]{t} \frac{\log^3 N}{N}. \quad \text{✶}$$

If we choose

$$N = 5,0246 t^{1/21} \log^{3/7} N$$

then the second and third terms are equal (except for the δ 's).

In order to round the calculations we adopt

$$(8.3) \quad N = 5t^{1/21}.$$

Then (8.1) becomes

$$(8.4) \quad \eta(t) = \sum_{q=1}^{\infty} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) \cdot t + \delta_1 \frac{5^6}{\pi^2} t^{2/7} + \delta_2 \frac{10240}{5} t^{2/7} \left(\log 5 + \frac{1}{21} \log t \right)^3.$$

Thus, in our evaluation we have included every possible pole at the primitive roots of unity.

Now

$$\begin{aligned} & \delta_1 \frac{5^6}{\pi^2} t^{2/7} + \delta_2 \frac{10240}{5} t^{2/7} \left(\log 5 + \frac{1}{21} \log t \right)^3 = \\ & = \delta_1 1583,14 t^{2/7} + \delta_2 2048 t^{2/7} \left(1,60943 + \frac{1}{21} \log t \right)^3 = \\ & = \delta_1 1583,14 t^{2/7} + \delta_3 2048 t^{2/7} \log^3 t = \delta_4 3631,14 t^{2/7} \log^3 t \end{aligned}$$

$-1 < \delta_i < 1$ for $t \geq 6$.

Hence (8.4) reduces to:

$$(8.5) \quad \eta(t) = \sum_{q=1}^{\infty} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) \cdot t + \delta_4 3631,14 t^{2/7} \log^3 t.$$

We know that due to the multiplicative character of its terms, the infinite series can be transformed into an infinite product, and therefore by [1] the final result is:

$$(8.6) \quad \sum_{q=1}^{\infty} \frac{\mu^2(q)}{\varphi^2(q)} Cq(t) = 2 \prod_{p=3}^{\infty} \left(1 - \frac{1}{(p-1)^2} \right) \prod_{p/t} \frac{p-1}{p-2} = 1,3203 \prod_{p/t} \frac{p-1}{p-2}.$$

Thus $\eta(t)$ consists of a first dominant discontinuous term

$$(8.7) \quad D(t) = 1,3203 \prod_{p/t} \frac{p-1}{p-2} t$$

and a second remainder term $C(t)$, for which we have the bound

$$(8.8) \quad C(t) = \delta_4 3631,14 t^{2/7} \log^3 t$$

with

$$(8.9) \quad \eta(t) = D(t) + C(t).$$

Recall that on account of (5.3) and (8.1) we have:

$$\eta(t) = \lim_{N \rightarrow \infty} L^{-1}\{B_N^2(s)\}.$$

9. Bounds for $\nu(t)$

From (7.4), letting $N \rightarrow \infty$ due to (7.1) and (3.9) we obtain:

$$\lim_{N \rightarrow \infty} f_N(t) = \frac{\nu(t+0) + \nu(t-0)}{2} = \int_0^1 u\eta(t-u)du + \int_1^2 (2-u)\eta(t-u)du.$$

Here we apply (8.9) in order to obtain:

$$(9.1) \quad \frac{\nu(t+0) + \nu(t-0)}{2} = D(t) \int_0^1 u \cdot du + D(t-1) \int_1^2 (2-u)du + \int_0^1 uC(t-u)du + \int_1^2 (2-u)C(t-u)du.$$

The last line can be evaluated by the mean value theorem of integral calculus, and we have

$$\begin{aligned} \frac{\nu(t+0) + \nu(t-0)}{2} &= \\ &= \frac{D(t) + D(t-1)}{2} + \delta_5 C(t - \delta_5) + (1 + \delta_6)C(t - 1 - \delta_6) = \\ &= \frac{D(t+0) + D(t-0)}{2} + 3\delta_7 C(t), \quad 0 < \delta_7 < 1, \quad 0 \leq \delta_5, \delta_6 < 1. \end{aligned}$$

Hence:

$$\nu(t) = D(t) + \frac{3}{2}\delta_7 C(t).$$

Appealing again to (8.7) and (8.8) for t even, we have:

$$\nu(t) = 1,303t \prod_{p|t} \frac{p-1}{p-2} t + 1,5 \delta_7 \delta_4 3631,14 t^{2/7} \log^3 t$$

from which follows the lower bound

$$(9.2) \quad \nu(t) > 1,303t - 5446,71 t^{2/7} \log^3 t$$

for every $t \geq 6$.

At this stage we can ask ourselves which is the greatest value that $\nu(t)$ can assume in the case that there is only one solution.

This solution can range from $p_1 = t - 3, p_2 = 3$ up to $p_1 = t/2, p_2 = t/2$. Hence $\max \nu(t) = \log^2 t/2$, or, better, since we must take into account the

order (because the solution $p_1 + p_2$ is counted differently from $p_2 + p_1$). Thus $\max \nu(t) = 2 \log^2 t/2$.

Consequently, if we know that

$$(9.3) \quad \nu(t) > 2 \log^2 t/2$$

we can be sure that there is at least one solution.

Replacing the lower bound (9.2) in (9.3) we have:

$$(9.4) \quad 1,303 t - 5446.71 t^{2/7} \log^3 t > 2 \log^2 t/2$$

which must hold if there is at least one solution.

Now, it is a numerical fact that (9.4) holds for $t > 10^{11}$. (Actually, if $t = 10^{11}$, we have $t^{2/7} = 1389,5046$, $\log t = 25,3284$, $\log^3 t = 16248,94324$.)

But the existence of the solutions has been proved up to $t = 4.10^{11}$ [11]. This is more than sufficient to prove the truth of the hypothesis.

10. Conclusion.

Using the same techniques, we can prove that **every** odd number is the sum of three primes. 75
^

Acknowledgement. By writing this paper I am honored to place on record my gratitude to one of the most distinguished mathematicians in this subject, Prof. Aldo Peretti. Without his generous help, advice and encouragement I could never have begun to work on this challenging problem.

The solution of Goldbach's conjecture in this paper is based almost entirely on his long and hard work about this problem, work which was published in some of his important papers in this matter.

In conclusion, the assistance and suggestions given by Prof. Peretti while writing this paper were valuable and very important.

To Prof. Aldo Peretti I address my warmest thanks.

References

- [1] HARDY, G., LITTLEWOOD, J., *Some problems of "Partitio numerorum"*, III: *On the expression of a number as a sum of primes*. Acta Math. 44 (1922), 1-70.
- [2] KRAWCYK, A., *On some property of the function $\varphi(n)$* . Acta Math. Univ. Comenianae (1980), 91-92.
- [3] LANDAU, E., *Sitzungber. Ak. Wiss. Wien (Math.)* 121, IIa, 1912, 2223.
- [4] LANDAU, E., *Vorlesungen über Zahlentheorie*, S. Hirzel (1927), Leipzig, vol.II.

- [5] PERETTI, A., *The solution of diophantine equations by means of the Laplace transform*. B.N.T. vol. II, april 1977, no.1, 1-12.
- [6] PERETTI, A., *The Goldbach problem (II)*. N.N.T.D.M. vol. 2 (1996), no.2, 53-68.
- [7] PERETTI, A., *The Goldbach problem (II)*. N.N.T.D.M. vol. (2) (1996), no.2, 1-26.
- [8] PERETTI, A., *The Goldbach's problem (III)*. N.N.T.D.M. vol. 2 (1997), no.4, 194-204.
- [9] PRUDNIKOV, A., BRYCHKOV, YU., MARICHEV, O., *Integrals and Series*. Vol.4. *Direct Laplace transforms*, vol. 5. *Inverse Laplace Transforms*. Gordon and Breach Science Publishers (1992).
- [10] REDMOND, D., *Number Theory*. Marcel Dekker, Inc. N.Y. (1996), Problem Set 4,2, p.202.
- [11] SINISALO, M., *Checking the Goldbach conjecture up to $4 \cdot 10^{11}$* Math. Comp. (1993), no.204, 931-934.
- [12] SNEDDON, I., *The use of integral transforms*. Tata Mc Graw Hill Publ. Co. Ltd. New Dehli (1974), §3.10, p.174.
- [13] VAUGHAN, R., *The Hardy-Littlewood method*. Cambridge University Press. 1982-1997, p.35, Formula 3, 23.

Accepted: 20.03.2000