

3

The Fibonacci Quarterly

THE OFFICIAL JOURNAL OF THE FIBONACCI ASSOCIATION

Veroff

VOLUME 21
NUMBER 4



NOVEMBER
1983

CONTENTS

Waiting Times and Generalized Fibonacci Sequences	<i>V.R.R. Uppuluri & S.A. Patil</i>	242
Powers of T and Soddy Circles	<i>John H. Selleck</i>	250
On Some Divisibility Properties of Fibonacci and Related Numbers	<i>Gerhard Rosenberger</i>	253
Letter to the Editor	<i>John Brillhart</i>	259
On the Solution of $(E^2 + (\lambda p - 2)E + (1 - \lambda p - \lambda^2 q))^n G_n = n^2$ by Expansions and Operators	<i>H.N. Malik</i>	260
On Fibonacci Numbers of the Form PX Where P is Prime	<i>Neville Robbins</i>	266
A General Method for Determining a Closed Formula for the Number of Partitions of the Integer n into m Positive Integers for Small Values of m	<i>W.J.A. Colman</i>	272
n -Dimensional Fibonacci Numbers and Their Applications	<i>Maluina Baica</i>	285
Counting the Profiles in Domino Tiling	<i>F.C. Wu</i>	302
The Fibonacci Sequences F_n Modulo L_n	<i>Karolyn A. Morgan</i>	304
Elementary Problems and Solutions	<i>Edited by A.P. Hillman, G.C. Padilla, & C.R. Wall</i>	306
Advanced Problems and Solutions	<i>Edited by Raymond E. Whitney</i>	312
Volume Index		319

♦♦♦♦

n -DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

MALVINA BAICA
Marshall University, Huntington WV 25701
(Submitted June 1982)

INTRODUCTION

In one of his papers [3] Bernstein investigated the $F(n)$ function. This function was derived from a special kind of numbers which could well be defined as 3-dimensional Fibonacci numbers. The original Fibonacci numbers should then be called 2-dimensional Fibonacci numbers. The present paper deals with n -dimensional Fibonacci numbers in a sense to be explained in the sequel. In a later paper [4] Bernstein derived an interesting identity that was based on 3-dimensional Fibonacci numbers. Also Carlitz in his paper [5] deals with this subject.

If we remember that the original Fibonacci numbers are generated by the formula

$$F(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-i}{i}, \quad n = 1, 2, \dots,$$

then the function

$$F(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^i \binom{n-2i}{i}$$

can be regarded as a generalization of the first, and the author thought that

$$F(n) = \sum_{i=0}^{\lfloor \frac{n}{k} \rfloor} (-1)^i \binom{n-ki}{i}, \quad k = 1, 2, \dots,$$

could serve as a k -dimensional generalization of the original Fibonacci numbers, but, regrettably, this consideration led nowhere. From the fact that the Fibonacci numbers are derived from the periodic expansion by the Euclidean algorithm of $\sqrt{5}$, there is opened a new horizon for the wanted generalization.

In a previous paper [1], the author had followed the ideas of Perron [9] and of Bernstein [4] and stated a general Algorithm that leads to an n -dimensional generalization of Fibonacci numbers.

In this paper, the author is introducing the GEA (Generalized Euclidean Algorithm) to investigate the various properties and applications of her k -dimensional Fibonacci numbers. It first turns out that these k -dimensional Fibonacci numbers are most useful for a good approximation of algebraic irrationals by rational integers. Further, the author proceeded to investigate higher-degree Diophantine equations and to state identities of a larger magnitude than those investigated before, in an explicit and simple form.

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

1. THE GEA

Let ω be the irrational

$$\begin{cases} \omega = \sqrt{D^n + 1}; n \geq 2, D \in N; \alpha^{(v)} = (\alpha_1^{(v)}(\omega), \dots, \alpha_{n-1}^{(v)}(\omega)), \\ \langle \alpha^{(v)} \rangle, \langle b^{(v)} \rangle \text{ sequences of the form } \alpha^{(v)}, v = 0, 1, \dots \end{cases} \quad (1.1)$$

The GEA of the fixed vector $\alpha^{(0)}$ is the sequence $\langle \alpha^{(v)} \rangle$ obtained by the recurrence formula

$$\begin{cases} \alpha^{(v+1)} = (\alpha_1^{(v)} - b_1^{(v)})^{-1} (\alpha_2^{(v)} - b_2^{(v)}, \dots, \alpha_{n-1}^{(v)} - b_{n-1}^{(v)}, 1) \\ b_i^{(v)} = \alpha_i^{(v)}(D); i = 1, \dots, n-1; v = 0, 1, \dots; \alpha_1^{(v)} \neq b_1^{(v)}. \end{cases} \quad (1.2)$$

The GEA of $\alpha^{(0)}$ is called purely periodic if there exists a number m such that

$$\alpha^{(0)} = \alpha^{(m)}; m \text{ is called the length of the primitive period} \quad (1.3)$$

The following formulas were proved in [2]. Let

$$\begin{cases} A_s^{(v+n)} = \sum_{k=0}^{n-1} b_k^{(v)} A_s^{(v+k)}; v = 0, 1, \dots \\ A_i^{(j)} = \delta_i^j; \delta_i^j \text{ the Kronecker delta, } i, j = 0, 1, \dots, n-1; s = 0, 1, \dots, n-1; \\ b_k^{(v)} = \alpha_k^{(v)}(D); k = 0, 1, \dots, n-1; \alpha_0^{(v)} = b_0^{(v)} = 1; \end{cases} \quad (1.4)$$

$A_s^{(v)}$ are called the matrices of GEA; then the three formulas hold:

$$\begin{vmatrix} A_0^{(v)} & A_0^{(v+1)} & \dots & A_0^{(v+n-1)} \\ A_1^{(v)} & A_1^{(v+1)} & \dots & A_1^{(v+n-1)} \\ \dots & \dots & \dots & \dots \\ A_{n-1}^{(v)} & A_{n-1}^{(v+1)} & \dots & A_{n-1}^{(v+n-1)} \end{vmatrix} = (-1)^{v(n-1)} \quad (1.5)$$

$$\alpha_s^{(0)} = \frac{\sum_{k=0}^{n-1} \alpha_k^{(v)} A_s^{(v+k)}}{\sum_{k=0}^{n-1} \alpha_k^{(v)} A_s^{(v+k)}}, v = 0, 1, \dots; s = 0, \dots, n-1. \quad (1.6)$$

$$\prod_{k=1}^v \alpha_{n-1}^{(k)} = \sum_{k=0}^{n-1} \alpha_k^{(v)} A_0^{(v+k)}. \quad (1.7)$$

Perron proved the following theorem which, under the conditions of the GEA ($D \geq 1$), becomes

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

Theorem 1

The GEA is convergent in the sense that

$$\left\{ \alpha_s^{(0)} = \frac{\lim_{\nu \rightarrow \infty} A_0^{(\nu)}}{\lim_{\nu \rightarrow \infty} A_3^{(\nu)}}, s = 1, \dots, n-1 \right. \quad (1.8)$$

$A_s^{(\nu)} : A_0^{(\nu)}$ is called the ν th convergent of GEA.

In [1], the author proved

Theorem 2

If the GEA of $\alpha^{(0)}$ is purely periodic with $m =$ length of the primitive period, then

$$\left\{ \begin{aligned} \prod_{k=0}^{m-1} \alpha_{n-1}^{(k)} &= \sum_{k=0}^{m-1} \alpha_k^{(m)} A_0^{(m+k)}, \\ &\text{is a unit in } \mathcal{Q}(\omega). \end{aligned} \right. \quad (1.9)$$

From (1.9) the formula follows, in virtue of (1.7),

$$\left(\prod_{k=0}^{m-1} \alpha_{n-1}^{(k)} \right)^{\nu} = \sum_{k=0}^{m-1} \alpha_k^{(\nu)} A_0^{(\nu m+k)}, \nu = 1, 2, \dots \quad (1.10)$$

2. A PERIODIC GEA

In this section, we construct a periodic GEA, with length of primitive period $m = 1$. The fixed vector $\alpha^{(0)}$ must be chosen accordingly, and this may look complicated at first. We prove

Theorem 3

The GEA of the fixed vector

$$\left\{ \begin{aligned} \alpha^{(0)} &= (\alpha_1^{(0)}, \alpha_2^{(0)}, \dots, \alpha_s^{(0)}, \dots, \alpha_{n-1}^{(0)}) \\ \alpha_s^{(0)} &= \sum_{i=0}^s \binom{n-s-1+i}{i} \omega^{s-i} D^i \\ &s = 1, \dots, n-1 \end{aligned} \right. \quad (2.1)$$

is purely periodic and the length of its primitive period $m = 1$.

Proof: We shall first need the formula

$$\sum_{i=0}^s \binom{n-s-1+i}{i} = \binom{n}{s}, \quad s = 1, \dots, n-1. \quad (2.2)$$

- n -DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

This is proved by induction. The proof is left to the reader. We have, from (2.1), the following components of $\alpha^{(0)}$ which we shall use later:

$$\alpha_1^{(0)} = w + (n-1)D; \alpha_{n-1}^{(0)} = \sum_{i=0}^{n-1} w^{n-1-i} D^i. \quad (2.3)$$

Since $w^n - D^n = 1$, we also have

$$\sum_{i=0}^{n-1} w^{n-1-i} D^i = (w - D)^{-1}. \quad (2.4)$$

The vectors $b_i^{(\nu)}$ ($i = 1, \dots, n-1; \nu = 0, 1, \dots$) obtained from $\alpha_i^{(\nu)}(w)$ by the defining rule (1.2) are called their corresponding companion vectors. We shall calculate the companion vector $b^{(0)}$ of $\alpha^{(0)}$ and have

$$b_s^{(0)} = \sum_{i=0}^s \binom{n-s-1+i}{i} D^{s-i} D^i = D^s \sum_{i=0}^s \binom{n-s-1+i}{i},$$

so that, by (2.2),

$$b_s^{(0)} = \binom{n}{s} D^s, \quad s = 1, 2, \dots, n-1. \quad (2.5)$$

Thus,

$$b^{(0)} = \left(\binom{n}{1} D, \binom{n}{2} D^2, \dots, \binom{n}{n-1} D^{n-1} \right).$$

We shall now calculate the vector $\alpha^{(1)}$. From (1.2), it follows that

$$\alpha^{(1)} = (\alpha_1^{(0)} - b_1^{(0)})^{-1} (\alpha_2^{(0)} - b_2^{(0)}, \dots, \alpha_{n-1}^{(0)} - b_{n-1}^{(0)}, 1). \quad (2.6)$$

From (2.3), (2.4), and (2.5), we obtain:

$$\begin{cases} \alpha_1^{(0)} - b_1^{(0)} = w + (n-1)D - \binom{n}{1} D = w - D, \\ \alpha_{n-1}^{(1)} = (w - D)^{-1} = \sum_{i=0}^{n-1} w^{n-1-i} D^i = \alpha_{n-1}^{(0)}. \end{cases} \quad (2.7)$$

We can prove the relation

$$(\alpha_s^{(0)} - b_s^{(0)}) (\alpha_1^{(0)} - b_1^{(0)})^{-1} = \alpha_{s-1}^{(1)}, \quad s = 2, \dots, n-1. \quad (2.8)$$

Since the proof is elementary, we leave it to the reader.

From (2.6), it follows that

$$\begin{cases} (\alpha^{(1)} = \alpha_1^{(0)}, \alpha_2^{(0)}, \dots, \alpha_{n-2}^{(0)}, \alpha_{n-1}^{(0)}) = \alpha^{(0)}, \\ \alpha^{(\nu)} = \alpha^{(0)}, \quad \nu = 1, 2, \dots \end{cases} \quad (2.9)$$

This proves Theorem 3.

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

3. EXPLICIT MATRICIANS $A_0^{(\nu+n)}$

We shall proceed to find an explicit formula for the "zero-degree matrices" $A_0^{(\nu+n)}$, $\nu = 0, 1, \dots$, and shall make use, for this purpose, of the defining formula (1.4), and the fact that the GEA is purely periodic with length of the primitive period $\bar{n} = 1$. Taking into account (2.5) and (2.9), we have

$$A_0^{(\nu+n)} = \sum_{s=0}^{n-1} \binom{n}{s} D^s A_0^{(\nu+s)}; \quad \nu = 0, 1, \dots \quad (3.1)$$

We shall now make use of Euler's generating function. We have

$$\begin{aligned} \sum_{i=0}^{\infty} A_0^{(i)} x^i &= x^0 A_0^{(0)} + \sum_{i=1}^{n-1} A_0^{(i)} x^i + \sum_{i=n}^{\infty} A_0^{(i)} x^i \\ &= 1 + \sum_{i=1}^{n-1} x^{i+n} \left(A_0^{(i)} + \binom{n}{1} D A_0^{(i+1)} + \binom{n}{2} D^2 A_0^{(i+2)} + \dots + \binom{n}{n-1} D^{n-1} A_0^{(i+n-1)} \right) \\ &= 1 + x^n \sum_{i=0}^{n-1} A_0^{(i)} x^i + x^{n-1} \sum_{i=0}^{n-1} \binom{n}{1} D A_0^{(i+1)} x^{i+1} + x^{n-2} \sum_{i=0}^{n-1} \binom{n}{2} D^2 A_0^{(i+2)} x^{i+2} \\ &\quad + \dots + x \sum_{i=0}^{n-1} \binom{n}{n-1} D^{n-1} A_0^{(i+n-1)} x^{i+n-1} \\ &= 1 + \left(x^n + \binom{n}{1} D x^{n-1} + \binom{n}{2} D^2 x^{n-2} + \dots + \binom{n}{n-1} D^{n-1} x \right) \sum_{i=0}^{n-1} A_0^{(i)} x^i \\ &\quad - \left(\binom{n}{1} D x^{n-1} + \binom{n}{2} D^2 x^{n-2} + \dots + \binom{n}{n-1} D^{n-1} x \right) = \sum_{i=0}^{n-1} A_0^{(i)} x^i. \end{aligned}$$

$$\begin{aligned} \left[1 - \left(x^n + \sum_{k=1}^{n-1} \binom{n}{k} D^k x^{n-k} \right) \right] \sum_{i=0}^{n-1} A_0^{(i)} x^i &= 1 - \sum_{k=1}^{n-1} \binom{n}{k} D^k x^{n-k}, \\ \sum_{i=0}^{n-1} A_0^{(i)} x^i &= \frac{1 - \left(x^n + \sum_{k=1}^{n-1} \binom{n}{k} D^k x^{n-k} \right) + x^n}{1 - \left(x^n + \sum_{k=1}^{n-1} \binom{n}{k} D^k x^{n-k} \right)} \\ &= \frac{x^n}{1 - \sum_{k=1}^{n-1} \binom{n}{k} D^k x^{n-k}} + 1, \end{aligned}$$

$$A_0^{(0)} + \sum_{i=1}^{n-1} A_0^{(i)} x^i = x^n \sum_{k=0}^{n-1} \left(\sum_{i=0}^{n-1} \binom{n}{k} D^k x^{n-k} \right)^i + 1.$$

$$x A_0^{(1)} + A_0^{(2)} x^2 + \dots + A_0^{(n-1)} x^{n-1} + \sum_{i=0}^{n-1} A_0^{(i)} x^i = x^n \sum_{k=0}^{n-1} \left(\sum_{i=0}^{n-1} \binom{n}{k} D^k x^{n-k} \right)^i,$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

For x sufficiently small. Thus, since $A_0^{(1)} = \dots = A_0^{(n-1)} = 0$, we have

$$\begin{aligned} \sum_{i=n}^{\infty} A_0^{(i)} x^i &= x^n \sum_{k=0}^{\infty} \left(\sum_{j=0}^{n-1} \binom{n}{k} D^k x^{n-k} \right)^n, \\ \sum_{i=0}^{\infty} A_0^{(n+i)} x^{n+i} &= x^n \sum_{k=0}^{\infty} \left(\sum_{j=0}^{n-1} \binom{n}{k} D^k x^{n-k} \right)^n, \\ \sum_{i=0}^{\infty} A_0^{(n+i)} x^i &= \sum_{k=0}^{\infty} \left(\sum_{j=0}^{n-1} \binom{n}{k} D^k x^{n-k} \right)^n, \end{aligned} \tag{3.2}$$

and comparing coefficients of powers x^v on both sides of (3.2), we obtain

$$A_0^{(v+n)} = \sum_{\substack{y_1 + y_2 + \dots + y_n = v \\ ny_1 + (n-1)y_2 + \dots + 2y_{n-1} + y_n = v}} \binom{v_1 + y_2 + \dots + y_n}{y_1, y_2, \dots, y_n} \prod_{k=0}^{n-1} \left(\binom{n}{k} D^k \right)^{y_{k+1}} \tag{3.3}$$

or

$$A_0^{(v+n)} = \sum_{\substack{y_1 + y_2 + \dots + y_n = v \\ \sum_{k=0}^{n-1} (n-k)y_{k+1} = v}} \binom{y_1 + y_2 + \dots + y_n}{y_1, y_2, \dots, y_n} D^{\sum_{k=0}^{n-1} y_{k+1}} \prod_{k=0}^{n-1} \left(\binom{n}{k} D^k \right)^{y_{k+1}}, \tag{3.4}$$

$$v = 0, 1, \dots$$

Formula (3.4) looks very complicated. $A_0^{(v+n)}$ can also be calculated by the recurrency relation (1.4). It is conjectured that it is easier to do so by formula (3.4), and would be a challenging computer problem.

4. MATRICIANS OF DEGREE $s, s = 1, 2, \dots, n-1$

In this section, we shall express "s-degree matricians,"

$$A_s^{(v)}, \quad s = 1, \dots, n-1,$$

by means of zero-degree matricians. This is not an easy task. Now we shall prove a very important theorem.

Theorem 4

The s-degree matricians are expressed through the zero-degree matricians by means of the relation

$$A_s^{(v+n-1)} = \sum_{k=0}^s \binom{n}{k} D^k A_0^{(v+n-s-k-1)}, \quad v = 0, 1, \dots; \tag{4.1}$$

$$s = 1, \dots, n-1.$$

Proof: From formula (1.6) it follows that

$$\alpha_s^{(0)} \sum_{k=0}^{n-1} \alpha_k^{(0)} A^{(v+k)} = \sum_{k=0}^{n-1} \alpha_k^{(0)} A_0^{(v+k)}, \quad s = 1, 2, \dots, n-1; \tag{4.2}$$

$$v = 0, 1, \dots$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

Or, writing a_i for $a_i^{(0)}$, $i = 0, \dots, n-1$, and substituting their values from (2.1), we obtain

$$\sum_{i=0}^s \binom{n-s-1+i}{i} \omega^{s-i} D^i \sum_{k=0}^{n-1} a_k A_0^{(v+k)} = \sum_{k=0}^{n-1} a_k A_s^{(v+k)}. \quad (4.3)$$

We shall now compare coefficients of ω^{n-1} on both sides of (4.3). The power of ω^{n-1} appears, on the right side only in

$$a_{n-1} = \omega^{n-1} + D \omega^{n-2} + \dots + D^{n-1},$$

and its coefficient is

$$A_0^{(v+n-1)}. \quad (4.4)$$

So the whole problem is to find the coefficient of ω^{n-1} on the left side, and this is the problem. We shall start with the first power of ω in a_s , which is ω^s (in the left side). Now in

$$\sum_{k=0}^{n-1} a_k A_0^{(v+k)}$$

we have to look for those a_k 's which have the powers ω^{n-s-1} ; this appears in

$$\begin{aligned} a_{n-s-1} & \left(\text{first term, coefficient} = A_0^{(v+n-s-1)} \right) \\ a_{n-s} & \left(\text{second term, coefficient} = \binom{s}{1} D A_0^{(v+n-s)} \right) \\ a_{n-s+1} & \left(\text{third term, coefficient} = \binom{s}{2} D^2 A_0^{(v+n-s+1)} \right) \\ & \vdots \\ a_{n-1} & \left((1+v)\text{th term, coefficient} = \binom{s}{s} D^s A_0^{(v+n-1)} \right). \end{aligned}$$

Thus, we have obtained the partial sum of coefficients of ω^{n-1} in the left side.

$$A_0^{(v+n-s-1)} + \binom{s}{1} D A_0^{(v+n-s)} + \binom{s}{2} D^2 A_0^{(v+n-s+1)} + \dots + \binom{s}{s} D^s A_0^{(v+n-1)}.$$

Now the next power of ω on the left side is ω^{s-1} with coefficient

$$\binom{n-s-1+1}{1} D = \binom{n-s}{1} D.$$

To obtain ω^{n-1} , ω^{s-1} must be multiplied by $v-s$, so we must start with the first term of a_{n-s} , the second term of a_{n-s+1} , ..., etc. Compared with the previous sum, s has to be replaced by $s-1$. The sum will then be multiplied by $\binom{n-s}{1} D$, and the number of summands will be smaller by one. We then obtain the partial sum:

$$\binom{n-s}{1} D \left[A_0^{(v+n-s)} + \binom{s-1}{1} D A_0^{(v+n-s+1)} + \binom{s-1}{2} D^2 A_0^{(v+n-s+2)} + \dots + \binom{s-1}{s-1} D^{s-1} A_0^{(v+n-1)} \right].$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

Thus the general term in the sum of coefficients of w^{n-1} on the left side of (4.3) which contains $D^k A_0^{(v+n-s-1+k)}$ as a constant factor has the form, adding up in (4.5) the column with this factor,

$$\sum_{i=0}^k \binom{n-s-1+i}{i} \binom{s-i}{k-i} D^k A_0^{(v+n-s-k+i)}. \quad (4.6)$$

The following formula is well known:

$$\sum_{i=0}^k \binom{n-s-1+i}{i} \binom{s-i}{k-i} = \binom{n}{k}, \quad (4.7)$$

which becomes formula (2.2) for $k = s$. Now, since in

$$a_s = \sum_{i=0}^s \binom{n-s-1+i}{i} w^{s-i} D^i,$$

the exponent of D sums from $i = 0$ to $i = s$, we have, finally,

$$A_s^{(v+n-1)} = \sum_{i=0}^s \binom{n}{i} D^i A_0^{(v+n-1-s+i)}$$

which is formula (4.1) and proves Theorem 4. From formula (4.1), we have the single cases

$$A_1^{(v+n-1)} = A_0^{(v+n-2)} + \binom{n}{1} D A_0^{(v+n-1)}, \quad (4.8)$$

and

$$A_{n-1}^{(v+n-1)} = A_0^{(v+n)}. \quad (4.9)$$

(4.9) is a very surprising relation and will be applied in the next section. Similarly,

$$A_2^{(v+n-1)} = A_0^{(v+n-3)} + \binom{n}{1} D A_0^{(v+n-2)} + \binom{n}{2} D^2 A_0^{(v+n-1)}, \text{ etc.} \quad (4.10)$$

5. APPROXIMATION OF IRRATIONALS BY RATIONALS

We shall investigate especially the case $D = 1$, but produce first formulas for any value of D . We obtain from (4.8) and (1.6),

$$\begin{aligned} \alpha_1^{(0)} &= \frac{\lim_{v \rightarrow \infty} A_1^{(v+n-1)}}{\lim_{v \rightarrow \infty} A_0^{(v+n-1)}} = \lim_{v \rightarrow \infty} \frac{A_0^{(v+n-2)} + n D A_0^{(v+n-1)}}{A_0^{(v+n-1)}} \\ &= \omega + (n-1)D = nD + \lim_{v \rightarrow \infty} \frac{A_0^{(v+n-2)}}{A_0^{(v+n-1)}}, \\ \omega &= D + \lim_{v \rightarrow \infty} \frac{A_0^{(v+n-2)}}{A_0^{(v+n-1)}} = D + \lim_{v \rightarrow \infty} \frac{A_0^{(v+n-1)}}{A_0^{(v+n)}}. \end{aligned} \quad (5.1)$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

For $D = 1$, $w = \sqrt[n]{2}$, and from (3.4) and (5.1) we obtain the approximation formula

$$\sqrt[n]{2} \approx \frac{\sum_{\sum_{i=0}^{n-1} y_{i+1} = v, i=0, \dots, n-1} \binom{y_1 + y_2 + \dots + y_n}{y_1, y_2, \dots, y_n} \prod_{k=0}^{n-1} b_{y_{k+1}}}{\sum_{\sum_{i=0}^{n-1} y_{i+1} = v+1, i=0, \dots, n-1} \binom{y_1 + y_2 + \dots + y_n}{y_1, y_2, \dots, y_n} \prod_{k=0}^{n-1} b_{y_{k+1}}}$$

$$b_k = \binom{n}{k}, \quad k = 0, \dots, n-1; \quad b_0 = 1.$$

The approximations are not very close, and we would have to continue a few steps further to get a closer approximation. Formula (4.9), surprisingly simple as it is, does not yield any news. It enables us to calculate w^{n-1} by means of the powers w_k , $k = 1, \dots, n-2$.

We have approximately, expanding $\sqrt[n]{2} = (1+1)^{1/n}$ by the binomial series,

$$\sqrt[n]{2} \approx 1 + \frac{1}{n}.$$

According to our approximation formula (5.1) with $D = 1$,

$$\sqrt[n]{2} = w = 1 + \frac{A_0^{(n)}}{A_0^{(n+1)}};$$

$$A_0^{(n+1)} = A_0^{(1)} + \binom{n}{1} A_0^{(2)} + \dots + \binom{n}{n-1} A_0^{(n)} = \binom{n}{n-1} A_0^{(n)} = n A_0^{(n)} = n,$$

since $A_0^{(n)} = A_0^{(0)} + A_0^{(1)} + \dots + \binom{n}{n-1} A_0^{(n-1)} = A_0^{(0)} = 1$, $\sqrt[n]{2} \approx 1 + \frac{1}{n}$, as should be.

6. DIOPHANTINE EQUATIONS

We shall construct two types of Diophantine equations of degree n in n unknowns and state their explicit solutions, which are infinite in number. We have from (1.5)

$$\begin{vmatrix} A_0^{(v+n)} & A_0^{(v+n+1)} & A_0^{(v+n+2)} & \dots & A_0^{(v+n+n-1)} \\ A_1^{(v+n)} & A_1^{(v+n+1)} & A_1^{(v+n+2)} & & A_1^{(v+n+n-1)} \\ \dots & \dots & \dots & \dots & \dots \\ A_{n-1}^{(v+n)} & A_{n-1}^{(v+n-1)} & A_{n-1}^{(v+n+2)} & & A_{n-1}^{(v+n+n-1)} \end{vmatrix} = (-1)^{(n-1)v}, \quad (6.1)$$

$$v = 0, 1, \dots$$

Substituting in (6.1) the values of $A_s^{(t)}$ from (4.1) we obtain, after simple row rearrangements,

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

$$\begin{vmatrix}
 A_0^{(\nu+n)} & A_0^{(\nu+n+1)} & A_0^{(\nu+n+2)} & \dots & A_0^{(\nu+n+n-1)} \\
 A_0^{(\nu+n-1)} & A_0^{(\nu+n)} & A_0^{(\nu+n+1)} & \dots & A_0^{(\nu+n+n-2)} \\
 A_0^{(\nu+n-2)} & A_0^{(\nu+n-1)} & A_0^{(\nu+n)} & \dots & A_0^{(\nu+n+n-3)} \\
 \dots & \dots & \dots & \dots & \dots \\
 A_0^{(\nu+3)} & A_0^{(\nu+4)} & A_0^{(\nu+5)} & \dots & A_0^{(\nu+n+2)} \\
 A_0^{(\nu+2)} & A_0^{(\nu+3)} & A_0^{(\nu+4)} & \dots & A_0^{(\nu+n+1)} \\
 A_0^{(\nu+1)} & A_0^{(\nu+2)} & A_0^{(\nu+3)} & \dots & A_0^{(\nu+n)}
 \end{vmatrix} = (-1)^{(n-1)\nu} \quad (6.3)$$

We introduce the notations

$$X_{\nu, k} = A_0^{(\nu+k)}, \quad k = 1, 2, \dots, n. \quad (6.3)$$

$$\begin{cases}
 A_0^{(\nu+k)} = A_0^{(\nu+k-n)} + b_1 A_0^{(\nu+k-n+1)} + b_2 A_0^{(\nu+k-n+2)} + \dots + b_{n-1} A_0^{(\nu+k-1)} \\
 b_k = \binom{n}{k} \sigma^k, \quad k = 0, 1, \dots, n-1, \quad \nu = 1, 2, \dots
 \end{cases} \quad (6.4)$$

We introduce these notations in (6.2) and then make the following manipulations in this determinant.

From the first row we subtract the b_1 multiple of the first row from below, then the b_2 multiple of the second row from below, ..., then the b_k th multiple of the k th row from below, $k = 1, \dots, n-1$.

Then (6.2) takes the form, in virtue of (6.4),

$$\begin{vmatrix}
 X_{\nu, n} - \sum_{k=1}^{n-1} b_k X_{\nu, k} & X_{\nu, 1} & X_{\nu, 2} & \dots & X_{\nu, n-1} \\
 A_0^{(\nu+n-1)} & A_0^{(\nu+n)} & A_0^{(\nu+n+1)} & \dots & A_0^{(\nu+n+n-2)} \\
 A_0^{(\nu+n-2)} & A_0^{(\nu+n-1)} & A_0^{(\nu+n)} & \dots & A_0^{(\nu+n+n-3)} \\
 \dots & \dots & \dots & \dots & \dots \\
 A_0^{(\nu+2)} & A_0^{(\nu+3)} & A_0^{(\nu+4)} & \dots & A_0^{(\nu+n+1)} \\
 X_{\nu, 1} & X_{\nu, 2} & X_{\nu, 3} & \dots & X_{\nu, n}
 \end{vmatrix} = (-1)^{(n-1)\nu} \quad (6.5)$$

We further subtract from the second row the b_2 multiple of the first row from below, the b_3 multiple of the second row from below, ..., the b_k multiple of the $(k-1)$ th row from below; the determinant (6.5) then takes the form ($k = 2, \dots, n-2$):

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

$$\begin{cases} X_{v,n} - \sum_{k=1}^{n-1} b_k X_{v,k} & X_{v,1} & & X_{v,2} & & X_{v,3} & & \dots & X_{v,n-1} \\ X_{v,n-1} - \sum_{k=1}^{n-2} b_{k+1} X_{v,k} & X_{v,n} - \sum_{k=1}^{n-1} b_{k+1} X_{v,k+1} & X_{v,1} + b_1 X_{v,2} & X_{v,2} + b_1 X_{v,3} & \dots & X_{v,n-1} + b_1 X_{v,n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ X_{v,1} & X_{v,2} & X_{v,3} & X_{v,4} & \dots & X_{v,n} \\ = & (-1)^{(n-1)v} \end{cases} \quad (6.6)$$

Continuing this process by another step, the third row of determinant (6.6) will have the form

$$X_{v,n-2} - \sum_{k=1}^{n-3} b_{k+2} X_{v,k} \quad X_{v,n-1} - \sum_{k=1}^{n-2} b_{k+2} X_{v,k+1} \quad X_{v,n} - \sum_{k=1}^{n-1} b_{k+2} X_{v,k+2} \\ X_{v,1} + b_1 X_{v,2} + b_2 X_{v,3} \quad X_{v,2} + b_1 X_{v,3} + b_2 X_{v,4} \quad \dots$$

$$X_{v,n-3} + b_1 X_{v,n-2} + b_2 X_{v,n-1}$$

Generally we subtract from the i th row in (6.2) the b_i multiple of the first row from below, then the b_{i+1} multiple of the second row from below, ..., the b_{n-1} multiple of the $(n-i)$ th row from below ($i = 1, \dots, n-1$). The reader can verify, that by these operations the determinant (6.2) transforms into one containing only the unknowns $X_{v,i}$ ($i = 1, \dots, n$), which yields the Diophantine equation of degree n in these unknowns.

7. MORE DIOPHANTINE EQUATIONS

The GEA of $a^{(0)}$ is purely periodic with length of the primitive period $m = 1$. Since

$$a_{n-1}^{(0)} = \sum_{i=0}^{n-1} \binom{n-1-i}{i} w^{n-1-i} D^i = \sum_{i=0}^{n-1} w^{n-1-i} D^i$$

we have by Theorem 2 and formula (1.10),

$$(w^{n-1} + Dw^{n-2} + \dots + D^{n-1})^v = \sum_{i=0}^{n-1} a_i^{(0)} A_0^{(v+i)}, \quad v = 1, 2, \dots \quad (7.1)$$

We find the norm of $(w^{n-1} + Dw^{n-2} + \dots + D^{n-1})^v$. We have

$$\begin{cases} D^n - w^n = -1, \\ D^n - w^n = -\sum_{k=0}^{n-1} (D - \rho_k w) = -N(D - w), \\ \rho_k = e^{2\pi i k/n}, \quad k = 0, 1, \dots, n-1. \end{cases} \quad (7.2)$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

But $\omega^{n-1} + D\omega^{n-2} + \dots + D^{n-1} = -(D - \omega)^{-1}$; hence,

$$N[(\omega^{n-1} + D\omega^{n-2} + \dots + D^{n-1})^v] = (-1)^{(n-1)v}, \quad \begin{matrix} v = 1, 2, \dots; \\ n = 2, 3, \dots \end{matrix} \quad (7.3)$$

We have

$$\begin{aligned} \sum_{i=0}^{n-1} a_i^{(0)} A_0^{(v+i)} &= A_0^{(v)} + \left[\omega + \binom{n-1}{1} D \right] A_0^{(v+1)} + \left[\omega^2 + \binom{n+2}{1} D\omega + \binom{n-1}{2} D^2 \right] A_0^{(v+2)} \\ &+ \left[\omega^3 + \binom{n-3}{1} \omega^2 D + \binom{n-2}{2} \omega D^2 + \binom{n-1}{3} D^3 \right] A_0^{(v+3)} + \dots \\ &+ \left[\omega^{n-1} + \binom{1}{1} \omega^{n-1} D + \binom{2}{2} \omega^{n-2} D^2 + \dots + \binom{n-1}{n-1} D^{n-1} \right] A_0^{(v+n-1)}. \end{aligned}$$

Denoting

$$\begin{cases} X_{v,k} = \sum_{s=0}^{n-1-k} \binom{n-1-k}{s} A_0^{(v+s+k)} D^s, \\ k = 0, 1, \dots, n-s. \end{cases} \quad (7.4)$$

This $X_{v,k}$ is not the $X_{v,k}$ from (6.4). We have from (7.1),

$$(\omega^{n-1} + D\omega^{n-2} + \dots + D^{n-1})^v = \sum_{k=0}^{n-1} X_{v,k} \omega^k = e^v, \quad e \text{ a unit.} \quad (7.5)$$

We shall find the field equation of

$$\sum_{k=0}^{n-1} X_{v,k} \omega^k.$$

The free member of it is the norm of e^v , and since e^v is a unit with the norm $(-1)^{(n-1)v}$, according to (7.3), we find easily, by known methods, that

$$\begin{vmatrix} X_{v,0} & X_{v,1} & X_{v,2} & \dots & X_{v,n-2} & X_{v,n-1} \\ mX_{v,n-1} & X_{v,0} & X_{v,1} & \dots & X_{v,n-3} & X_{v,n-2} \\ mX_{v,n-2} & mX_{v,n-1} & X_{v,0} & \dots & X_{v,n-4} & X_{v,n-3} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ mX_{v,2} & mX_{v,3} & mX_{v,4} & \dots & X_{v,0} & X_{v,1} \\ mX_{v,1} & mX_{v,2} & mX_{v,3} & \dots & mX_{v,n-1} & X_{v,1} \end{vmatrix} = (-1)^{(n-1)v} \quad (7.6)$$

It is not difficult to see that, in the case $n = 2m + 1$ ($m = 1, 2, \dots$), the highest powers of the n unknowns of the discriminant (7.6) as

$$X_{v,0}^m, mX_{v,1}^m, m^2X_{v,2}^m, \dots, m^{m-1}X_{v,m-1}^m$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

while the last unknown, $X_{v, n-1}$ does not have the exponent n , but a smaller one. In the case $n = 2m$ ($m = 1, 2, \dots$) these $n - 1$ powers are the same, but with alternating signs, viz.,

$$X_{v,0}^n, -mX_{v,1}^n, +m^2X_{v,2}^n, \dots$$

In the case $n = 2$, the expanded discriminant (7.6) had the form

$$X_v^2 - mY_v^2 = \pm 1,$$

and in the case $n = 3$, it had the form

$$X^3 + mY^3 + m^2Z^3 - 3mXYZ = 1.$$

The first is Pell's equation.

8. IDENTITIES AND UNITS

We return to formulas (7.4) and (7.5), and have

$$\left\{ \begin{array}{l} X_{nv,k} = \sum_{s=0}^{n-1-k} \binom{n-1-k}{s} A_0^{(vn+s+k)} D^s \\ k = 0, 1, \dots, n-1 \\ (w^{n-1} + Dw^{n-2} + \dots + D^{n-1}) = \sum_{k=0}^{n-1} X_{nv,k} w^k. \end{array} \right. \quad (8.1)$$

We compare powers of w^k ($k = 0, 1, \dots, n-1$) on both sides of (8.1) and take into consideration that $w^{nv} = m^v = (D^n + 1)^v$. We have, looking for the rational part of the right side, $k = 0$, and the value of the right side equals $X_{nv,0}$, and by (7.4);

$$X_{nv,0} = \sum_{s=0}^{n-1} \binom{n-1}{s} A_0^{(nv+s)} D^s, \quad v = 0, 1, \dots \quad (8.2)$$

On the left side, we have to look for the coefficients of w^n . Since the highest power in the expression

$$(w^{n-1} + Dw^{n-2} + \dots + D^{n-1})^{nv}$$

is $n(n-1)v$, we have the expression

$$\sum_{\substack{\sum_{t=1}^{n-1} (n-t)y_t = nv \\ \sum_{t=1}^{n-1} y_{t+1} = n(n-1)v - nv, \quad \sigma = 0, 1, \dots, (n-1)v}} \binom{y_1 + y_2 + \dots + y_n}{y_1, y_2, \dots, y_n} w^{\sum_{t=1}^{n-1} (n-t)y_t} D^{\sum_{t=1}^{n-1} y_{t+1}} = X_{nv}, \quad (8.3)$$

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

We want to obtain in this way the rational part of

$$(w^{n-1} + Dw^{n-2} + \dots + D^{n-1})^{nv}.$$

At the same time

$$\sum_{i=1}^{n-1} iy_{i+1}$$

is the sum of the exponents of the powers of y_{i+1} ($i = 1, \dots, n-1$). Since in every summand of

$$w^{n-1} + Dw^{n-2} + \dots + D^{n-1}$$

the sum of the exponents of $D^i w^{n-1-i}$ ($i = 0, 1, \dots, n-1$) is $n-1$, and the highest exponent in the expansion is $n(n-1)v$, we have that

$$\sum_{i=1}^{n-1} (n-i)y_i - \sum_{i=1}^{n-1} iy_{i+1} = n(n-1)v,$$

which explains the left side of (8.3). We further have

$$\sum_{i=1}^{n-1} [(n-i)y_i + iy_{i+1}] = n(n-1)v,$$

so that

$$y_1 + y_2 + \dots + y_n = nv. \quad (8.4)$$

Now, taking into account that the exponent of w under the summation sign in (8.3) equals sn , $w^{sn} = m^s$, and $D^s = m-1$, formula (8.3) takes the form

$$\left\{ \begin{aligned} & \sum_{i=1}^{n-1} \sum_{\substack{y_1 + y_2 + \dots + y_n = nv \\ y_i \geq 0}} \binom{nv}{y_1, y_2, \dots, y_n} m^s (m-1)^{(n-1)v-s} = X_{nv,0} \\ & = \sum_{k=0}^{n-1} \binom{n-1}{k} D^k A_0^{(nv+k)}, \\ & s = 0, 1, \dots, (n-1)v; \nu = 0, 1, \dots \\ & y_1 + y_2 + \dots + y_n = nv \end{aligned} \right. \quad (8.5)$$

(8.5) is an interesting combinatorial identity.

From (8.1), $n-1$ more identities can be obtained by comparing the coefficients of the powers w^i , $i = 1, \dots, n-1$, on both sides of (8.1). The identities have a somewhat complicated form; however, they will express the coefficients of w^t , $t = 1, \dots, n-1$, in the expansion of

$$(w^{n-1} + Dw^{n-2} + \dots + D^{n-1})^{nv},$$

with $w^n = m = D^n + 1$;

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

$$\left\{ \begin{aligned} & \sum_{i=1}^{n-1} (n-i)y_i = sn + t \leq n(n-1)v \\ & \sum_{i=1}^{n-1} iy_{i+1} = n(n-1)v - (sn + t) \\ & = \sum_{j=0}^{n-1-t} \binom{n-1-t}{j} K_0^{(nv+j+t)} D^j \\ & j = 0, 1, \dots, (n-1)v - 1; t = 1, \dots, n-1. \end{aligned} \right. \quad (8.6)$$

We wish to explain the appearance of the factor D^{n-t} under the summation sign on the left side of (8.6). The power of D in the expansion of

$$(D^{n-1} + D^{n-2} + \dots + D^{n-1})^{vn}$$

equals

$$\begin{aligned} \sum_{i=1}^{n-1} iy_{i+1} &= n(n-1)v - (sn + t) \\ &= n(n-1)v - sn - n + (n-t) \\ &= n[(n-1)v - s - 1] + n - t. \end{aligned}$$

Thus, the power of D equals

$$(D^n)^{n(n-1)v-s-1} \cdot D^{n-t}, \text{ with } D^n = m - 1.$$

The power of w is

$$\sum_{i=1}^{n-1} (n-i)y_i = sn + t = (w^n)^s w^t = m^s w^t,$$

so m^s is the coefficient of w^t as desired.

REFERENCES

1. M. Baica. "An Algorithm in a Complex Field (And Its Applications to the Calculation of Units). Accepted for publication by the *Pacific Journal of Mathematics*.
2. L. Bernstein. "The Jacobi-Perron Algorithm, Its Theory and Application." *Lecture Notes in Mathematics*, No. 207 (Springer-Verlag, 1971), I-IV, 1-160.
3. L. Bernstein. "Zeroes of the Function $F(n) = \sum_{i=0}^{n-2i} (-1)^i \binom{n-2i}{i}$." *Journal of Number Theory* 6, No. 4 (1974):264-70.
4. L. Bernstein. "A Formula for Fibonacci Numbers from a New Approach to Generalized Fibonacci Numbers." *The Fibonacci Quarterly* 14, No. 4 (1976): 358-68.

n-DIMENSIONAL FIBONACCI NUMBERS AND THEIR APPLICATIONS

5. L. Carlitz. "Some Combinatorial Identities of Bernstein." *SIAM J. Math. Analysis* 9 (1978):65-75.
6. L. Carlitz. "Recurrences of the Third Order and Related Combinatorial Identities." *The Fibonacci Quarterly* 16, No. 1 (1978):11-18.
7. P. Erdos. "Arithmetical Properties of Polynomials." *J. London Math. Soc.* 28 (1953):416-25.
8. V. E. Hoggatt, Jr. *Fibonacci and Lucas Numbers*. Boston: Houghton-Mifflin Co., 1969, pp. 1-91.
9. O. Perron. "Grundlagen fuer Eine Theorie des Jacobischen Kettenbruchalgorithmus." *Math. Ann.* 64 (1970):1-76.
10. H. Zassenhaus. "On the Units of Order." *Journal of Algebra* 20 (1972):368-95.

♦♦♦♦♦

♦♦♦♦

VOLUME INDEX

- AINSWORTH, O. R. (coauthor J. Neggers). "A Family of Polynomials and Powers of the Secant," 21(2):132-38.
- BAICA, Malvina. " n -Dimensional Fibonacci Numbers and Their Applications," 21(4):285-301.
- BANGE, David W. (coauthor A. E. Barkauskas). "Fibonacci Graceful Graphs," 21(3):174-88.
- BARKAUSKAS, Anthony E. (coauthor D. W. Bange). "Fibonacci Graceful Graphs," 21(3):174-88.
- BRILLHART, John. "Letter to the Editor," 21(4):259.
- COHEN, M. E. (coauthor D. L. Hudson). "On Exponential Series Expansions and Convolutions," 21(2):111-17.
- COLMAN, W. J. A. "A General Method for Determining a Closed Formula for the Number of Partitions of the Integer n into m Positive Integers for Small Values of m ," 21(4):272-84.
- DARBRO, Wesley A. (coauthor G. von Tiesenhausen). "Sequences Generated by Self-Replicating Systems," 21(2):97-106.
- DE BOUVÈRE, Karel L. (coauthor Regina E. Lathrop). "Injectivity of Extended Generalized Fibonacci Sequences," 21(1):37-52.
- DEO, Narsingh (coauthor M. Quinn). "Pascal Graphs and Their Properties," 21(3):203-14.
- EGECIOGLU, Ömer. "The Parity of the Catalan Numbers via Lattice Paths," 21(1):65-66.
- EHRHART, E. "Associated Hyperbolic and Fibonacci Identities," 21(2):87-96.
- ERLEBACH, Lee (coauthor W. Y. Vélez). "Equiprobability in the Fibonacci Sequence," 21(3):189-91.
- GARCIA, P. G. (coauthor S. Ligh). "A Generalization of Euler's ϕ -Function," 21(1):26-28.
- GEORGHIOU, C. (coauthor A. N. Philippou). "Harmonic Sums and the Zeta Function," 21(1):29-36.
- GIRSE, Robert D. "A Note on Fibonacci Cubature," 21(2):129-31.
- GODSIL, Christopher D. (coauthor R. Razen). "A Property of Fibonacci and Tribonacci Numbers," 21(1):13-17.
- HALE, David R. "A Variant of Nim and a Function Defined by Fibonacci Representation," 21(2):139-42.
- HILLMAN, A. P., Ed. Elementary Problems and Solutions, 21(1):67-73; 21(2):147-52; 21(3):230-35; 21(4):306-11.
- HOCK, J. L. (coauthor R. B. McQuistan). "The Occupational Degeneracy for λ -Bell Particles on a Saturated $\lambda \times n$ Lattice Space," 21(3):196-202.
- HORIBE, Yasuichi. "Notes on Fibonacci Trees and Their Optimality," 21(2):118-28.
- HUDSON, D. L. (coauthor M. E. Cohen). "On Exponential Series Expansions and Convolutions," 21(2):111-17.
- KIMBERLING, Clark. "One-Free Zeckendorf Sums," 21(1):53-57.
- KIRSCHENHOFER, Peter (coauthors H. Prodinger & R. F. Tichy). "Fibonacci Numbers of Graphs: II," 21(3):219-29.
- LATHROP, Regina E. (coauthor K. L. de Bouvère). "Injectivity of Extended Generalized Fibonacci Sequences," 21(1):37-52.
- LAWTON, Wayne M. "Kronecker's Theorem and Rational Approximation of Algebraic Numbers," 21(2):143-46.
- LEHMER, D. H. (coauthor Emma Lehmer). "Properties of Polynomials Having Fibonacci Numbers for Coefficients," 21(1):62-64.
- LEHMER, Emma (coauthor D. H. Lehmer). "Properties of Polynomials Having Fibonacci Numbers for Coefficients," 21(1):62-64.
- LIGH, Steve (coauthor P. G. Garcia). "A Generalization of Euler's ϕ -Function," 21(1):26-28.
- MALIK, H. N. "On the Solution of $\{E^2 + (\lambda p - 2)E + (1 - \lambda p - \lambda^2 q)\}^m G_n = n^k$ by Expansions and Operators," 21(4):260-65.